

Corporate Medical Policy

Urinary Tumor Markers for Bladder Cancer AHS – G2125

File Name: urinary_tumor_markers_for_bladder_cancer
Origination: 1/1/2019
Last Review: 5/2024

Description of Procedure or Service

A. Definitions

Bladder cancer is defined as a malignancy that develops from the tissues of the bladder. It is the most common cancer of the urinary system. The cancer typically arises from the urothelium, although it may originate in other locations such as the ureter or urethra (Lerner, 2023).

Tumor biomarkers are proteins detected in the blood, urine, or other body fluids that are produced by the tumor itself or in response to it. Urinary tumor markers may be used to help detect, diagnose, and manage some types of cancer including bladder cancer (Hottinger & Hormigo, 2011).

Related Policies

Serum Tumor Markers for Malignancies AHS – G2124
Liquid Biopsy AHS – G2054

B. Background

Each year in the United States, the American Cancer Society estimates there are about 82,290 new cases of bladder cancer and about 16,710 deaths from bladder cancer (ACS, 2023). Bladder cancer is the sixth most common cancer in the United States, affects men four times more frequently than women, and is typically diagnosed in individuals above the age of 40, with 73 the median age at diagnosis (DeGeorge et al., 2017; NCCN, 2023a). Bladder cancer risk factors include smoking, a family history of the disease, pelvic radiation, obesity, diabetes, and chronic infection of the urinary tract.

Bladder cancer commonly presents as painless hematuria (blood in urine) and may be gross (visible) or microscopic. Gross hematuria tends to increase the likelihood of bladder cancer, but hematuria as a whole may be transient or due to non-cancer related causes (Perazalla, 2021). Other common symptoms of bladder cancer include pain or irritative and obstructive voiding symptoms such as urge incontinence, dysuria, straining, or nocturia. These symptoms are often mistaken for another condition such as kidney stones, can be temporary, and are not necessarily specific for bladder cancer (Lotan, 2022). In fact, hematuria is the most common symptom of bladder cancer, but a study reported a 13% prevalence rate of bladder cancer out of 6728 patients with hematuria (DeGeorge et al., 2017; Sutton et al., 2018). Approximately 70%-75% of patients present with superficial tumors (50 – 70% of which can recur but are usually not life threatening), and 25%-30% present as invasive tumors with a high risk of metastasis (Chou & Dana, 2010; Kaufman et al., 2009).

Cystoscopy (white light) is the gold standard for a diagnosis of bladder cancer. This procedure involves a bladder examination and urine sample for cytology. Any lesions are observed and recorded. Cystoscopy does not detect all malignancies or visualize the upper urinary tract.

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Furthermore, although cystoscopy is minimally invasive, it may be uncomfortable and promote anxiety, which can lead to suboptimal compliance with management recommendations. Fluorescent cystoscopy is somewhat more efficient at detecting tumors than white light cystoscopy; although, it comes with its own set of issues such as higher false-positive rates and costs (Lotan & Choueiri, 2022; Mitra et al., 2023). Cytology, or the analysis of cells in urine, is often completed in addition to cystoscopy analysis.

Although cystoscopy has long been the gold standard for a diagnosis of bladder cancer, its high cost and unpleasant burden has led to the search for a non-invasive test that can match the high specificities and sensitivities set by cystoscopy. Urinary biomarkers including “Cell-free proteins and peptides, exosomes, cell-free DNA, methylated DNA and DNA mutations, circulating tumor cells, miRNA, lncRNA, rtRNA and mRNAs” have now been identified for bladder cancer diagnostic purposes (Lopez-Beltran et al., 2019). Urine is exposed to urothelial tissue in many different locations, and therefore has the potential to contain several biomarkers associated with cancer. Validation of these biomarkers could lessen the use of cystoscopy as well as increase the overall sensitivity for bladder cancer identification (D'Costa et al., 2016). However, because of the lower disease prevalence in a screening population, even in those at increased risk, the use of biomarkers for screening is not cost effective or recommended (Lotan et al., 2009). Despite the promise of urine biomarkers, cystoscopy remains the procedure of choice both for initial diagnosis and for surveillance in previously treated patients.

Epigenetic changes may also play an important role in bladder cancer tumorigenesis. These changes are becoming more prevalent as identification rates increase due to improvements in high-throughput DNA sequencing technologies. Epigenetic changes can “regulate [the] gene expression outcome without changing the underlying DNA sequence” with alterations based on DNA methylation, nucleosome positioning, microRNA regulation and histone modifications. All these epigenetic-based changes are distorted in each human cancer type. “A substantial portion (76%) of all primary bladder tumors displays mutations in at least one chromatin regulatory gene. These mutations cause epigenetic dysregulation in bladder cancers” (Li et al., 2016).

Numerous other urinary biomarkers have been proposed as contributors to management of bladder cancer.

Other nuclear matrix proteins aside from NMP22 have been investigated. NMP52, BLCA-4, and BLCA-1 have all been studied as potential markers. Initial data for these markers appears promising, but most likely requires further evaluation (Mitra et al., 2023).

Cytokeratins, protein components of the cell structure, have also been identified as possible markers. Cytokeratins (“CK”), -8, -18, -19, and -20 have been considered for use in bladder cancer evaluation. However, further data is needed (Mitra et al., 2023).

Other markers that have been considered as potential indicators of bladder cancer include the following:

Telomerase is an enzyme that adds telomeres to the ends of chromosomes. This enzyme is only expressed in proliferating cells such as cancer cells, thereby lending credence to its use as a cancer marker. Despite its high sensitivity, its clinical application is limited, as the current assay used to detect telomerase is “significantly” affected by sample collection and processing (Mitra, et al., 2023).

Hyaluronic acid is a polysaccharide that promotes tumor progression and metastasis. It is cleaved by *hyaluronidase*, which creates smaller fragments of the polysaccharide that further promote tumor angiogenesis. This pair of markers has been found to detect low-grade and low-stage disease with higher sensitivities than other markers, but requires further data for evaluation (Mitra et al., 2023).

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Fibrin degradation products may also be useful in detection of cancer. High levels of vascular endothelial growth factor can increase the permeability of surrounding cellular structures, which cause serum proteins to “leak”. These proteins are eventually degraded to fibrin, and then to fibrin degradation products (Mitra et al., 2023).

Survivin is an apoptosis inhibitor. Survivin is frequently elevated in cancers, but virtually undetectable in normal tissues. However, no commercial assays for Survivin exist as of time of writing (Mitra et al., 2023).

Finally, *miRNA* markers have been considered for use in bladder cancer management. These markers are small sequences of non-coding RNA that contribute to gene expression regulation. MiRNAs-126, -200c, -143, and -222 have all been considered to have “promising” results (Mitra et al., 2023).

Proprietary Testing

The two most studied urinary biomarkers are bladder tumor antigen (BTA) and nuclear matrix protein 22 (NMP22). The BTA test is designed to detect complement factor H-related protein (hCFHrp) which is elevated in cancer cells. This test is available in both a quantitative and qualitative version, and its manufacturer-recommended cut-off is 14U/MI (Mahnert et al., 1999; Mitra et al. 2023). The BTA stat® test and the BTA TRAK® test are available from Polymedco and measure qualitative and quantitative detection of bladder tumor-associated antigen, respectively. Similarly, the NMP22 test is designed to detect a protein that is more highly available in cancer cells than normal cells. In this case, cancer cells release more NMP22 into the urine following apoptosis than normal cells do. The NMP22 tests are also available in a quantitative and qualitative version, and its FDA-approved cut-off is 10U/MI (Grossman et al., 2005; Mitra et al., 2023; Zuiverloon et al., 2017). A number of proprietary tests exist revolving around one of these two biomarkers; these tests include Abbott’s “Alere NMP22 BladderCheck” and Quest’s Bladder Tumor Antigen DetectR (Abbott, 2023; Quest, 2020).

The FDA has approved two additional tests for urinary biomarkers. One is UroVysion, which is designed to detect chromosomal alterations that are distinctive of bladder cancer. This test is a fluorescent in situ hybridization (FISH) assay that uses DNA probes to detect alterations (such as aneuploidies) on chromosomes 3, 7, and 17 or loss of the 9p21 locus. The second test is known as ImmunoCyt (or uCyt+) that uses a similar fluorescent technique to detect certain glycoproteins that are expressed solely on cancerous cells (Mitra et al., 2023).

Recently, Pangea Laboratory has created a laboratory developed test termed Bladder CARE™ which measures the methylation status of specific DNA biomarkers in urine for the detection of bladder cancer via an at-home collection kit. This non-invasive test has not been approved by the FDA, is purported to be more cost-effective, and uses an epigenetic-based detection approach. Specifically, the methylation of bladder cancer DNA biomarkers are measured (Pangea, 2020). As little as 5 ng of urine DNA from a 100 MI urine sample is required, and it has a limit detection of 0.1% leading to the identification of a single cancerous cell in a sample of 1,000 normal cells (Pangea, 2020). The authors claim that Bladder CARE™ has a sensitivity of 94% and specificity of 86%, allowing for the identification of 88% of low-grade bladder cancer cases; these results are based on a study completed by Pangea Laboratory and Zymo Research which analyzes urine samples from 182 patients (97 with bladder cancer and 85 healthy controls) (Pangea, 2019).

Another test, termed the Bladder EpiCheck test, has been developed by the Israeli company Nucleix. This non-invasive epigenetic urine test helps to detect bladder cancer with a panel of 15 DNA methylation biomarkers. Nucleix reports a sensitivity of 92%, a specificity of 88% and a negative predictive value of 99% for the Bladder EpiCheck test; these results are based on a multi-center clinical study with 353 bladder cancer patients (Nucleix, 2015). Similar results have

Urinary Tumor Markers for Bladder Cancer AHS – G2125

been reported by D'Andrea et al. (2019). However, this test is not available in the United States (Nucleix, 2015).

Another test, termed “UBC® Rapid” has been developed by the Swedish company ODL Biotech. This point-of-care test measures soluble fragments of cytokeratins 8 and 18 in urine samples. The test can produce results within 10 minutes and may be tested with hematuria-containing samples. UBC® Rapid is the only quantitative point of care test platform for urine-based detection of bladder cancer. (AroCell, 2023). Ecke et al. (2018) performed a validation of this test, which encompassed 242 patients with bladder cancer (134 non-muscle-invasive low-grade tumors, 48 non-muscle-invasive high-grade tumors, 60 muscle-invasive high-grade tumors), 62 patients with non-evidence of disease [NED], and 226 healthy controls. The authors found a sensitivity of 38.8% for non-muscle-invasive low-grade bladder cancer, 75% for non-muscle-invasive high-grade bladder cancer and 68.3% for muscle-invasive high-grade bladder cancer. Specificity over the entire cohort was 93.8% (Ecke et al., 2018).

The URO17 assay by Protean Biodiagnostics, an immunohistochemistry-based test that detects the presence of the oncoprotein keratin 17 in bladder cancer and urogenital cancer. Unlike other urine-based test URO17 can detect patients with visible or invisible hematuria, which allows for early diagnosis. URO17 can also detect recurrent bladder cancer in patients under surveillance for relapse (NICE, 2023). The test has 100% sensitivity and 96% specificity for detecting bladder cancer from urine samples (Protean Biodiagnostics, 2021).

Nonagen Bioscience released Oncuria, an in-vitro multiplex immunoassay, which detects protein biomarkers associated with bladder cancer in the urine. This non-invasive test detects ten proteins from a single urine sample in patients with hematuria with suspicion of bladder cancer. Biomarker levels are combined in a weighted algorithm to aid in the prediction of responding to Bacillus Calmette-Guerin (BCG) therapy in patients with intermediate to high-risk, early-stage bladder cancer (Nonagen Bioscience, 2022).

The Xpert® Bladder Cancer Monitor can be used as a diagnostic in a population of patients with a history of non-muscle invasive bladder cancer (NMIBC). The test was designed for use in follow-up of patients undergoing routine surveillance. Pichler et al. (2018) enrolled 140 patients with a history of NMIBC and the patients underwent urine cytology using the Paris classification system. Urinary specimens were also analyzed with PCR using the Xpert® BC monitor, which measures five target mRNAs (ABL1, CRH, IGF2, UPK1B, and ANXA10). The overall sensitivity of the Xpert® BC Monitor was 0.84 with an NPV of 0.93. The authors write that this was “significantly superior” to the sensitivity of bladder washing cytology (0.33 and 0.76; $P < 0.001$). Another subgroup analysis confirmed the sensitivity as compared to barbotage cytology (Pichler et al., 2018).

D'Elia et al. (2021) also performed a study tracking follow-up and diagnostic utility of the Xpert® BC for patients with a history of NMIBC. This prospective study was done using 1015 samples from a group of 416 patients. Patients had a urinary cytology, the Xpert® Bladder Cancer monitor test, and cystoscopy. If the cystoscopy was positive, a transurethral resection of the bladder was completed. The Xpert® BC test identified 168 recurrent tumors: 126 were low-grade and 42 were high-grade; the overall sensitivity was 17.9% for cytology, 52.4% for the Xpert® BC test and 54.2% for the two tests combined. Overall specificity was 98.5% for cytology, 78.4% for the Xpert® BC test, and 78.2% for the two tests combined (D'Elia et al., 2021).

Analytical Validity

Piao et al. (2019) have developed a way to differentiate patients with bladder cancer from patients with a nonmalignant hematuria without bladder cancer by measuring urinary cell-free microRNA expression. This study shows that the non-invasive measurement of urinary

Urinary Tumor Markers for Bladder Cancer AHS – G2125

microRNA-6124 and microRNA-4511 can be used as a diagnostic tool with a sensitivity of >90% (Piao et al., 2019). This testing method will help to reduce the number of unnecessary cystoscopies in patients with hematuria that are being evaluated for bladder cancer.

The performance of an epigenetic-based bladder cancer detection tool has been evaluated by Fantony et al. (2017); the urine-based TWIST1/NID2 methylation assay has been analyzed for the detection of urothelial carcinoma via the addition of urine cytology. This multi-institutional study analyzed data from 172 patients. The authors note that “The AUC [area under the curve] for cytology alone with equivocal cytologies positive was 0.704 and improved to 0.773 with the addition of the DNA methylation assay ($p < 0.001$)” (Fantony et al., 2017). The authors conclude by stating that this TWIST1/NID2 methylation assay is a sensitive diagnostic tool that adds value to urine cytology for the detection of urothelial carcinoma, which is the most common type of bladder cancer.

Soubra and Risk (2015) found the sensitivity of fluorescent cystoscopy to be 0.92 and the sensitivity of white light cystoscopy to be 0.71; the specificity of fluorescent cystoscopy was lower at 0.57, and the specificity of white light cystoscopy was identified at 0.72. Furthermore, fluorescent cystoscopy’s sensitivity for carcinoma in situ (which is difficult to visualize) was measured at 0.924, while white light cystoscopy’s sensitivity for carcinoma in situ was much lower at 0.605, but these differences tended to decrease on higher grade lesions (Soubra & Risk, 2015). Cytology is also a common analytic technique in addition to cystoscopy. Its overall sensitivity is low at 0.34 and its sensitivity for grade 1 and 2 tumors is even lower at 0.12 and 0.26, respectively (Lotan & Roehrborn, 2003).

Breen et al. (2015) compared the sensitivity and specificity values of four diagnostic tests (cytology, NMP22, UroVysion, and CxBladder); CxBladder was found to have the highest sensitivity at 74% and cytology was identified with the highest specificity at 95%. The authors report comparable sensitivity values for cytology, NMP22, and UroVysion at 46%, 45.9% and 47.7% respectively (Breen et al., 2015). It is important to note that even though CxBladder is reported to have the highest sensitivity, the specificity (81.7%) is the lowest; the other tests were reported to have superior specificities with NMP22 at 88%, and UroVysion at 87.7% (Breen et al., 2015).

Sathianathen et al. (2018) published a study focusing on biomarkers in patients presenting with hematuria. This study encompassed BTA, NMP22, FISH, and uCyt+, as well as a fifth biomarker known as AssureMDx. Sensitivities ranged from 0.67 (BTA) to 0.95 (AssureMDx, second highest was uCyt+ at 0.83) while specificities ranged from 0.68 (BTA) to 0.93 (quantitative NMP22). However, this data is consistent with the previously published meta-analysis that covered all settings, not just hematuria (Chou et al., 2015). Cytology was also found to have superior specificity to all studied biomarkers; although, biomarkers tended to have better sensitivity. The authors concluded that, due to the high heterogeneity and small sample size, more studies were needed to validate biomarkers to replace diagnostic evaluation of hematuria (Sathianathen et al., 2018).

Although many studies emphasize the high validity of biomarkers such as NMP22 and BTA, these studies often have a large proportion of high-grade tumors which inflate the specificity and sensitivity; hence, the problem of identifying low-grade cancers remains. There may be changes at the genetic level in a low-grade cancer, but the proteins tested in the urine may still be relatively normal (D’Costa et al., 2016). Another issue is the conflicting results for the validity of the biomarkers. For example, the sensitivity of the quantitative NMP22 test has been found to range from as low as 0.26 to 1.00 with its specificity ranging from 0.49 to 0.98. Similarly, the BTA STAT test’s sensitivity and specificity have been found to range from 0.29 to 0.91 and from 0.54 to 0.86 respectively (Zuiverloon et al., 2017). For comparison, a study found the sensitivity and specificity of flexible cystoscopy (out of 778 hematuria patients) to be 0.98 and 0.938, respectively (Sutton et al., 2018).

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Dudley et al. (2019) have developed a novel high-throughput sequencing method that uses urine derived tumor DNA (utDNA) known as utDNA CAPP-Seq (Ucapp-Seq) to detect bladder cancer. This technique was used to analyze samples from 118 patients with early-stage bladder cancer and 67 healthy adults. “We detected utDNA pretreatment in 93% of cases using a tumor mutation-informed approach and in 84% when blinded to tumor mutation status, with 96% to 100% specificity” (Dudley et al., 2019). These results show that utDNA can be used to diagnose early-stage bladder cancer with high sensitivity and specificity.

Hirasawa et al. (2021) studied the diagnostic performance of Oncuria™, a multiplex immunoassay urinalysis test for bladder cancer. Urine samples from 362 subjects with suspicion of bladder cancer were measured using Oncuria™ for ten biomarkers (A1AT, APOE, ANG, CA9, IL8, MMP9, MMP10, PAI1, SDC1 and VEGFA). Results of the test were confirmed by cystoscopy and tissue biopsy. “The Oncuria™ test achieved a strong overall diagnostic performance, achieving an overall AUC of 0.95, sensitivity and specificity values of 93% and 93%, respectively, and a negative predictive value (NPV) and positive predictive value (PPV) of 99% and 65%, respectively. The Oncuria™ test shows promise for clinical application in the non-invasive diagnosis and surveillance bladder cancer, and potentially for screening at-risk, asymptomatic individuals” (Hirasawa et al., 2021).

Clinical Utility and Validity

A meta-analysis of 57 studies detailed the accuracy of several biomarkers for the diagnosis and surveillance of bladder cancer. These included the six FDA-approved tests (quantitative and qualitative NMP22, quantitative and qualitative BTA, FISH, and uCyt+) as well as a laboratory developed test that does not require FDA approval termed CxBladder. Sensitivities ranged from 0.57 (qualitative NMP22) to 0.82 (CxBladder); however, the CxBladder cohort was only comprised of one study. The specificities ranged from 0.74 (quantitative BTA) to 0.88 (qualitative NMP22). Sensitivity increased as a tumor progressed (higher grade or stage) with low accuracy for lower stage or grade tumors. A cytologic evaluation performed with a biomarker assessment increased sensitivity as well but missed about 10% of cases. Ultimately, the authors concluded that urinary biomarkers reported many false-positive results and failed to identify a large percentage of patients with bladder cancer (Chou et al., 2015). The authors also noted that this was the first study which focused on the measurement of clinical outcomes based on urinary biomarkers.

The ideal marker will be “easier, better, faster, and cheaper” (Schmitz-Dräger et al., 2015). Overall, although there have been numerous promising studies for the clinical utility of these urinary biomarkers, the biomarkers do not yet measure up to the standards set by cystoscopy as the primary method of diagnosis. Most of the biomarkers are yet to be well-validated and the ones that are, such as NMP22 and BTA, fall short of cystoscopy’s standards (D’Costa et al., 2016). Furthermore, because of the lower disease prevalence in a screening population, even in those at increased risk, the use of biomarkers for screening is not cost effective or recommended (Lotan et al., 2009). Although the cost of tests is non-clinical, it is still a crucial issue; the BTA and NMP22 tests are relatively inexpensive at \$25 but ImmunoCyt costs around \$80 and the CxBladder and UroVysion cost \$325 and \$800, respectively (Zuiverloon et al., 2017). For comparison, a cystoscopy cost around \$210 in 2016, and a cystoscopy with a biopsy cost about \$370 (Halpern et al., 2017). These biomarkers to date have not been highly recommended within any clinical guidelines. Therefore, the authors concluded that biomarkers have not had significant effect on clinical decision-making (Schmitz-Dräger et al., 2015). The majority of studies performed on these biomarkers did not focus on their ability to predict the course of cancer (D’Costa et al., 2016) but some biomarkers may play a role in the diagnosis or surveillance of bladder cancer in the future (Schmitz-Dräger et al., 2015).

An in-depth health technology assessment (HTA) of Cxbladder test was performed by Landaas et al. (2020) integrating clinical data and real-world usage scenarios to highlight the test’s sensitivity and specificity. Data from a vendor-funded study showed sensitivity of 91% and specificity of 60% for Cxbladder; another study indicated a sensitivity of 0.82 and specificity of

Urinary Tumor Markers for Bladder Cancer AHS – G2125

.85. The authors also noted an Agency for Healthcare Research and Quality (AHRQ)-funded systematic review by Chou et al. (2015) highlighting the high false-positive rate and poor accuracy of Cxbladder for low-stage and low-grade tumors. The AHRQ concluded that urinary biomarkers like Cxbladder would miss a substantial portion of bladder cancer cases and tests were subject to false positive results (Chou et al., 2015).

A follow-up pilot study by Landaas et al. (2020) was initiated at UW Medicine to analyze the best use-case scenario for Cxbladder. The pilot study involved patients with a history of urothelial carcinoma, comparing those tests with Cxbladder (group 1) to a control group (group 2). Group 1 patients underwent various follow-up tests including urine cytologies, cystoscopies, and biopsies, with recurrence detected in two out of six patients within the study period. Group 2, without Cxbladder testing, had three out of six patients with detected recurrence. The study essentially found no significant differences in follow-up tests between the two groups. These findings underscore the complexities of adopting new molecular diagnostic tests like Cxbladder on a system-wide basis. However, the study did find that Cxbladder testing was beneficial for a specific patient profile: those with normal cystoscopy results and atypical cytology. In such cases, Cxbladder testing led to fewer follow-up procedures (cystoscopies, cytologies, and biopsies) while still detecting a similar proportion of bladder cancer recurrences as standard procedures within the year. In conclusion, Cxbladder appears most suitable for those undergoing surveillance for bladder cancer recurrence, particularly those with normal cystoscopy and atypical cytology (Landaas et al., 2020).

The majority of studies performed on these biomarkers did not focus on their ability to predict the course of cancer (D'Costa et al., 2016) but some biomarkers may play a role in the diagnosis or surveillance of bladder cancer in the future (Schmitz-Dräger et al., 2015). Even this may be a difficult barrier to cross; Meleth et al. (2014) prepared an assessment for the Agency for Healthcare Research and Quality that stated “although UroVysion is marketed as a diagnostic rather than a prognostic test, limited evidence from two small studies (total n=168) supported associations between test result and prognosis for risk of recurrence” (Meleth et al., 2014). The authors went on to note that no studies that established clinical utility were found.

D'Andrea et al. (2019) analyzed 357 urine samples from patients at five different centers under surveillance for non-muscle-invasive bladder cancer to investigate the clinical utility of the Bladder EpiCheck™ non-invasive urine test. A specificity of 88% was identified with this test, a negative predictive value of 94.4% for the detection of any cancer, and a negative predictive value of 99.3% for the detection of high grade cancer; the use of the Bladder EpiCheck™ test helped to improve the cancer recurrence predictive value by a difference of 16-22% (D'Andrea et al., 2019). This high-performing diagnostic test may help in the surveillance of non-muscle-invasive bladder cancer.

Tan et al. (2018) completed a systematic review to identify the diagnostic sensitivity and specificity of urinary biomarkers for the diagnosis of bladder cancer. The authors report that multi-target biomarker panels were more accurate than single biomarker targets, and that both the sensitivity and specificity of urinary biomarkers were higher in primary diagnostic scenarios compared to patients under surveillance (Tan et al., 2018). The authors note that “few biomarkers achieve a high sensitivity and negative predictive value,” with single biomarkers reporting a sensitivity of 2-94% and specificity of 46-100%, and multi-target biomarkers reporting a sensitivity of 24-100% and specificity of 48-100% (Tan et al., 2018).

Mossanen et al. (2019) performed a cost analysis to characterize the costs of managing non-muscle-invasive bladder cancer (NMIBC). The authors created a Markov model with four health states: no evidence of disease, recurrence, progression and cystectomy, and death. Patients were stratified into three risk categories of low, intermediate, and high. The authors found that “cumulative costs of care over a 5-year period were \$52,125 for low-risk, \$146,250 for intermediate-risk, and \$366,143 for high-risk NMIBC.” The authors identified that the primary driver of cost was “progression to muscle-invasive disease requiring definitive therapy”, which

Urinary Tumor Markers for Bladder Cancer AHS – G2125

was found to contribute 81% and 92% to overall cost for intermediate and high-risk disease, respectively. Progression of disease was found to contribute 71% to overall cost for low-risk disease. The authors concluded that although protracted surveillance cystoscopy does contribute to management cost, progression of disease was the dominant factor in increasing cost of care (Mossanen et al., 2019).

Vasdev et al. (2021) studied the role of URO17™ biomarker in the diagnosis of bladder or urothelial cancer in new hematuria patients. Urine samples from 71 subjects were stained using the URO17™ immunobiomarker and results were compared to the biopsy and histology. URO17™ was shown to have an overall sensitivity of 100%, specificity of 92.6%, positive predictive value of 0.957, and negative predictive value of 1. URO17™ investigation was positive in every case of urothelial malignancy. According to the authors, URO17™ test can help improve “diagnostic capabilities in primary care, reduce the number of referrals to Urology department, and reduce the number of unnecessary invasive procedures for new patients with a suspected urinary bladder cancer” (Vasdev et al., 2021).

C. State and Federal Regulations, as applicable

Food and Drug Administration (FDA)

On April 16, 1997, the FDA approved the Bard BTA stat™ Test, created by Bard Diagnostic Sciences Inc. From the FDA site: “the BTA stat test is an in vitro diagnostic immunoassay indicated for the qualitative detection of bladder tumor associated antigen in urine of persons diagnosed with bladder cancer. This test is indicated for use as an aid in the management of bladder cancer patients in conjunction with cystoscopy.”

On April 15, 1998, the FDA approved the BTA TRAK™ Test, created by Bard Diagnostic Sciences Inc. From the FDA site: “the BTA TRAK test is an in vitro diagnostic immunoassay indicated for the quantitative detection of bladder tumor associated antigen in human urine. This test is indicated for use as an aid in the management of bladder cancer patients in conjunction with cystoscopy.”

On July 2, 1996, the FDA approved the MATRITECH NMP22™ TEST KIT, created by Alere Scarborough Inc. From the FDA site: “The Matritech NMP22 Test Kit is an enzyme immunoassay (EIA) for the in vitro quantitative determination of nuclear matrix protein NMP22 in stabilized voided urine.”

On July 30, 2002, the FDA approved the NMP22 BladderChek, created by Matritech Inc. From the FDA site: “The Matritech NMP22 BladderChek Test is indicated for professional and prescription home use as an aid in monitoring bladder cancer patients, in conjunction with standard diagnostic procedures.” This assay is qualitative.

On January 24, 2005, the FDA approved the UROVYSION BLADDER CANCER KIT. From the FDA site: “The UroVysion Bladder Cancer Kit (UroVysion Kit) is designed to detect aneuploidy for chromosomes 3, 7, 17, and loss of the 9p21 locus via fluorescence in situ hybridization (FISH) in urine specimens from persons with hematuria suspected of having bladder cancer.”

On February 23, 2000, the FDA approved the ImmunoCyt, created by Diagnocure Inc. From the FDA site: “ImmunoCyt is a qualitative direct immunofluorescence assay intended for use in conjunction with cytology to increase overall sensitivity for the detection of tumor cells exfoliated in the urine of patients previously diagnosed with bladder cancer. ImmunoCyt is indicated for use as an aid in the management of bladder cancer in conjunction with urinary cytology and cystoscopy” (FDA, 2018).

Urinary Tumor Markers for Bladder Cancer AHS – G2125

All of the FDA-approved tests apart from ImmunoCyt are approved for both diagnosis and surveillance of bladder cancer whereas ImmunoCyt is only approved for surveillance (Darwiche et al., 2015).

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

*****Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

Policy

BCBSNC will provide coverage for Urinary Tumor Markers for Bladder Cancer when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Urinary Tumor Markers for Bladder Cancer is covered

1. Reimbursement is allowed for urinary biomarkers (bladder tumor antigen (BTA) test, nuclear matrix protein (NMP22) test, or fluorescence in situ hybridization (FISH) UroVysion Bladder Cancer test):
 - a. as an adjunct in the diagnostic exclusion of bladder cancer for individuals who have an atypical or equivocal cytology
 - b. as an adjunct in the monitoring of high-risk, non-muscle invasive bladder cancer
2. Reimbursement is allowed for the use of fluorescence immunocytology (ImmunoCyt/uCyt) as an adjunct to cystoscopy or cytology in the monitoring of individuals with bladder cancer.

When Urinary Tumor Markers for Bladder Cancer is not covered

1. Reimbursement is not allowed for urinary biomarkers (bladder tumor antigen (BTA) test, nuclear matrix protein (NMP22) test, or fluorescence in situ hybridization (FISH) UroVysion Bladder Cancer test) when used for screening of bladder cancer in asymptomatic individuals, evaluation of hematuria, diagnosing bladder cancer in symptomatic individuals, and all other indications.
2. Reimbursement is not allowed for the use of fluorescence immunocytology (ImmunoCyt/uCyt) in the evaluation of hematuria, diagnosing bladder cancer in symptomatic individuals, or screening for bladder cancer in asymptomatic persons.
3. Any other urinary tumor markers for bladder cancer not mentioned above do not meet coverage criteria.

Table of Terminology

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Term	Definition
AACC	American Association for Clinical Chemistry
ACS	American Cancer Society
AHRQ	Agency for Healthcare Research and Quality
AMH	Asymptomatic microhematuria
ASCO	American Society of Clinical Oncology
ASTRO	American Society for Radiation Oncology
AUA	American Urological Association
AUC	Area under the curve
BC	Bladder cancer
BCG	Bacillus 10urvivin-guerin
BLCA-1	Bacillus collagen-like protein of anthracis
BLCA-4	Bacillus collagen-like protein of anthracis
BTA	Bladder tumor antigen
CDC	Centers For Disease Control and Prevention
CFHrp	Complement factor h-related protein
CIS	Carcinoma in situ
CK	Cytokeratins
CLIA '88	Clinical Laboratory Improvement Amendments of 1988
CMS	Centers for Medicare and Medicaid Services
CXCR2	C-X-C motif chemokine receptor 2
DNA	Deoxyribonucleic acid
EAU	European Association of Urology
EIA	Enzyme immunoassay
FDA	United States Food and Drug Administration
FISH	Fluorescence in situ hybridization
hCFHrp	Complement factor h-related protein
HTA	Health technology assessment
ICUD-SIU	International Consultation on Urological Diseases & Société Internationale d'Urologie
LDTs	Laboratory-developed tests
MH	Microhematuria
MRI	Magnetic resonance imaging
NACB	National Academy of Clinical Biochemistry Laboratory Medicine
NCCN	National Comprehensive Cancer Network
NCI	National Cancer Institute
NED	Non-evidence of disease
NID2	Nidogen 2
NMIBC	Non-muscle invasive bladder cancer

Urinary Tumor Markers for Bladder Cancer AHS – G2125

NMP22	Nuclear matrix protein 22
NMP52	Nuclear matrix protein 52
PCR	Polymerase chain reaction
RCTs	Randomized controlled trials
SUFU	Society of Urodynamics, Female Pelvic Medicine & Urogenital Reconstruction
SUO	Society of Urologic Oncology
TWIST1	Twist-related protein 1
uCyt+	<i>ImmunoCyt</i> test
USPSTF	U.S. Preventive Services Task Force
UT	Urine derived tumor
utDNA	Urine derived tumor deoxyribonucleic acid

Policy Guidelines

Guidelines and Recommendations

National Comprehensive Cancer Network (NCCN)

The NCCN has stated that “Urine molecular tests for urothelial tumor markers are now available. Many of these tests have a better sensitivity for detecting bladder cancer than urinary cytology, but specificity is lower. Considering this, evaluation of urinary urothelial tumor markers may be considered during surveillance of high-risk non-muscle-invasive bladder cancer. However, it remains unclear whether these tests offer additional information that is useful for detection and management of non-muscle-invasive bladder tumors. Therefore, the panel considers this to be a category 2B recommendation” (NCCN, 2023b)

American Urological Association (AUA)

The AUA’s guidelines on the diagnosis, evaluation and follow-up of asymptomatic microhematuria (AMH) in adults do not recommend use of urine markers (NMP22, BTA-stat, UroVysion) as part of routine evaluation (Davis et al., 2012).

The AUA and Society of Urodynamics, Female Pelvic Medicine & Urogenital Reconstruction (SUFU) published a guideline on microhematuria in 2020. In it, they remark that “Clinicians should not use urine cytology or urine-based tumor markers in the initial evaluation of patients with microhematuria”, stating that “insufficient evidence exists that routine use would improve detection of bladder cancer.” However, the guideline states that “Clinicians may obtain urine cytology for patients with persistent microhematuria after a negative workup who have irritative voiding symptoms or risk factors for carcinoma in situ.” Overall, the guideline states that “the panel does not recommend using urine cytology or urine-based tumor markers in the initial evaluation of MH [microhematuria] because, to date, markers have not demonstrated incrementally additive information to cystoscopy in the MH population, not have they been found to be of sufficient predictive value to obviate cystoscopy” (Barocas et al., 2020).

The AUA and Society of Urologic Oncology (SUO) joint guidelines on Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer (NMIBC) do not recommend using urinary biomarkers to replace cystoscopy when monitoring NMIBC (grade B), although a clinician can use biomarkers to evaluate a patient’s response to Bacillus Calmette-Guerin (BCG) therapy or a

Urinary Tumor Markers for Bladder Cancer AHS – G2125

separate cytology such as FISH or ImmunoCyt. However, a urinary biomarker should not be used for monitoring a patient with a normal cystoscopy and a history of low-risk cancer (Chang et al., 2020). This 2016 guideline was amended in 2020, but no relevant changes were identified.

The 2021 American Urologic Association (AUA) annual meeting included a guideline amendment update for non-muscle invasive bladder cancer (NMIBC) and muscle-invasive bladder cancer (MIBC) to the 2020 guidelines. According to the update, a clinician should not use urinary biomarkers in place of cystoscopy. “In a patient with a history of low-risk cancer and a normal cystoscopy, a clinician should not routinely use a urinary biomarker or cytology during surveillance. In a patient with NMIBC, a clinician may use biomarkers to assess response to intravesical BCG (UroVysion® FISH) and adjudicate equivocal cytology (UroVysion® FISH and ImmunoCyt™)” (AUA/SUO, 2020). The panel does acknowledge the uptake of Cxbladder in clinical practice; however, there is a lack of high quality evidence in the potential replacement of cystoscopy with Cxbladder (AUA, 2021).

Similarly, the joint guidelines between the AUA, the SUO, the American Society of Clinical Oncology (ASCO), and the American Society for Radiation Oncology (ASTRO) regarding non-metastatic muscle-invasive bladder cancer note that molecular biomarkers may be important for staging cancer and deciding a course of treatment soon. Nevertheless, at this time the biomarkers have not been properly validated (Chang et al., 2017).

U.S. Preventive Services Task Force (USPSTF)

The USPSTF concluded in 2011 that there was insufficient evidence to evaluate screening for bladder cancer in asymptomatic adults, assigning a grade I to this recommendation. Since then, there have been no further guidelines published on this topic by the USPSTF (Moyer, 2011).

In 2021, the USPSTF published the following statement regarding bladder cancer screening in adults: “Literature scans conducted in November 2021 in the MEDLINE and PubMed databases and the Cochrane Library showed a lack of new evidence to support an updated systematic review on the topic at this time (USPSTF, 2021).

3rd International Consultation on Urological Diseases & Société Internationale d’Urologie (ICUD-SIU)

With a level of evidence of 3 and a grade of “B”, the ICUD-SIU recommends, “examination of urine cytology must be a part of the expectant management or active surveillance protocol.” Concerning the surveillance strategies for NMIBC, “Surveillance strategies following a negative 3 months surveillance cystoscopy should be: (1) for low-risk disease, cystoscopy 6–9 months later and annually thereafter; consider cessation following five recurrence-free years. No upper tract imaging necessary unless hematuria present; (2) for intermediate risk, cystoscopy with cytology every 3–6 months for 2 years; then every 6–12 months during years 3 and 4; then annually for lifetime. Upper tract imaging every 1–2 years; (3) for high risk, cystoscopy with cytology every 3 months for 2 years; then every 6 months during years 3 and 4; then annually for lifetime [Level of evidence: 3; Grade C]” (Monteiro et al., 2018).

National Cancer Institute (NCI)

In the 2022 update to the NCI’s Bladder and Other Urothelial Cancers Screening (PDQ®)—Health Professional Version, the NCI states that “There is inadequate evidence to determine whether screening for bladder and other urothelial cancers has an impact on mortality... Based on fair evidence, screening for bladder and other urothelial cancers would result in unnecessary diagnostic procedures with attendant morbidity” (NCI, 2022) .

European Association of Urology (EAU)

Urinary Tumor Markers for Bladder Cancer AHS – G2125

The EAU has published guidelines on non-muscle-invasive bladder cancer (NIBC).

In 2022, the EAU concluded that “Cystoscopy is necessary for the diagnosis of bladder cancer” and that “Urinary cytology has high sensitivity in high-grade tumours including carcinoma in situ.” The EAU remarks that “There is no known urinary marker specific for the diagnosis of invasive BC [bladder cancer]” (Witjes et al., 2022).

An update to guidelines on non-muscle-invasive bladder cancer (NIBC) was published in 2022. The EAU concluded that urinary molecular marker tests cannot replace cystoscopy in routine practice, “but the knowledge of positive test results (microsatellite analysis) can improve the quality of follow-up cystoscopy.” Diagnosis ultimately depends on “cystoscopy examination of the bladder and histological evaluation of sampled tissue” (Babjuk et al., 2022).

An update to the EAU guidelines was published in 2023. In it, the EAU commented on urinary molecular marker tests, “None of these markers have been accepted as routine practice by any clinical guidelines for diagnosis or follow-up.” However, they remarked that “promising urinary biomarkers, assessing multiple targets, have been tested in prospective multicentre studies. Four of the promising and commercially available urine biomarkers, Cxbladder, ADX-Bladder, Xpert Bladder and EpiCheck, although not tested in RCTs, have such high sensitivities and negative predictive values in the referenced studies for high grade disease that these biomarkers may approach the sensitivity of cystoscopy. These 4 tests might be used to replace and/or postpone cystoscopy as they may identify the rare HG recurrences occurring in low/intermediate NMIBC” (EAU, 2023).

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable codes: 86294, 86316, 86386, 88120, 88121, 88346, 88350, 0012M, 0013M, 0363U, 0365U, 0366U, 0367U, 0420U

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

Abbott. (2023). ALERE NMP22® BLADDERCHEK®. Retrieved 12/18/2023 from <https://www.globalpointofcare.abbott/en/product-details/nmp22-bladderchek.html>

ACS. (2023). Key Statistics for Bladder Cancer. <https://www.cancer.org/cancer/bladder-cancer/about/key-statistics.html>

AroCell. (2023). Bladder Cancer Detection. Retrieved 12/18/2023 from <https://arocell.com/products/oncology/ubc-rapid/>

AUA. (2021). AUA Guideline Amendment: Non-Muscle Invasive Bladder Cancer/Muscle Invasive Bladder Cancer. <https://www.urotoday.com/conference-highlights/aua-2021-program/aua-2021-bladder-cancer/131965-aua-2021-aua-guideline-amendment-non-muscle-invasive-bladder-cancer-muscle-invasive-bladder-cancer.html>

Urinary Tumor Markers for Bladder Cancer AHS – G2125

AUA/SUO. (2020). Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Joint Guideline (2020). <https://www.auanet.org/guidelines/guidelines/bladder-cancer-non-muscle-invasive-guideline>

Babjuk, M., Burger, M., Capoun, O., Cohen, D., Compérat, E. M., Dominguez Escrig, J. L., Gontero, P., Liedberg, F., Masson-Lecomte, A., Mostafid, A. H., Palou, J., van Rhijn, B. W. G., Rouprêt, M., Shariat, S. F., Seisen, T., Soukup, V., & Sylvester, R. J. (2022). European Association of Urology Guidelines on Non-muscle-invasive Bladder Cancer (Ta, T1, and Carcinoma in Situ). *European Urology*, 81(1), 75-94. <https://doi.org/10.1016/j.eururo.2021.08.010>

Barocas, D. A., Boorjian, S. A., Alvarez, R. D., Downs, T. M., Gross, C. P., Hamilton, B. D., Kobashi, K. C., Lipman, R. R., Lotan, Y., Ng, C. K., Nielsen, M. E., Peterson, A. C., Raman, J. D., Smith-Bindman, R., & Souter, L. H. (2020). Microhematuria: AUA/SUFU Guideline. *J Urol*, 204(4), 778-786. <https://doi.org/10.1097/ju.0000000000001297>

Breen, V., Kasabov, N., Kamat, A. M., Jacobson, E., Suttie, J. M., O'Sullivan, P. J., Kavalieris, L., & Darling, D. G. (2015). A holistic comparative analysis of diagnostic tests for urothelial carcinoma: a study of Cxbladder Detect, UroVysion® FISH, NMP22® and cytology based on imputation of multiple datasets. *BMC medical research methodology*, 15, 45-45. <https://doi.org/10.1186/s12874-015-0036-8>

Chang, S. S., Bochner, B. H., Chou, R., Dreicer, R., Kamat, A. M., Lerner, S. P., Lotan, Y., Meeks, J. J., Michalski, J. M., Morgan, T. M., Quale, D. Z., Rosenberg, J. E., Zietman, A. L., & Holzbeierlein, J. M. (2017). Treatment of Non-Metastatic Muscle-Invasive Bladder Cancer: AUA/ASCO/ASTRO/SUO Guideline. *J Urol*, 198(3), 552-559. <https://doi.org/10.1016/j.juro.2017.04.086>

Chang, S. S., Boorjian, S. A., Chou, R., Clark, P. E., Daneshmand, S., Konety, B. R., Pruthi, R., Quale, D. Z., Ritch, C. R., Seigne, J. D., Skinner, E. C., Smith, N. D., & McKiernan, J. M. (2020). Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Guideline. *J Urol*, 196(4), 1021-1029. <https://doi.org/10.1016/j.juro.2016.06.049>

Chou, R., & Dana, T. (2010). Screening adults for bladder cancer: A review of the evidence for the u.s. preventive services task force. *Annals of Internal Medicine*, 153(7), 461-468. <https://doi.org/10.7326/0003-4819-153-7-201010050-00009>

Chou, R., Gore, J. L., Buckley, D., Fu, R., Gustafson, K., Griffin, J. C., Grusing, S., & Selph, S. (2015). Urinary Biomarkers for Diagnosis of Bladder Cancer: A Systematic Review and Meta-analysis. *Ann Intern Med*, 163(12), 922-931. <https://doi.org/10.7326/m15-0997>

D'Andrea, D., Soria, F., Zehetmayer, S., Gust, K. M., Korn, S., Witjes, J. A., & Shariat, S. F. (2019). Diagnostic accuracy, clinical utility and influence on decision-making of a methylation urine biomarker test in the surveillance of non-muscle-invasive bladder cancer. *BJU Int*, 123(6), 959-967. <https://doi.org/10.1111/bju.14673>

D'Costa, J. J., Goldsmith, J. C., Wilson, J. S., Bryan, R. T., & Ward, D. G. (2016). A Systematic Review of the Diagnostic and Prognostic Value of Urinary Protein Biomarkers in Urothelial Bladder Cancer. *Bladder Cancer*, 2(3), 301-317. <https://doi.org/10.3233/blc-160054>

D'Elia, C., Folchini, D. M., Mian, C., Hanspeter, E., Schwienbacher, C., Spedicato, G. A., Pycha, S., Vjaters, E., Degener, S., Kafka, M., Pycha, A., & Trenti, E. (2021). Diagnostic value of Xpert® Bladder Cancer Monitor in the follow-up of patients affected by non-muscle invasive bladder cancer: an update. *Ther Adv Urol*, 13, 1756287221997183. <https://doi.org/10.1177/1756287221997183>

Urinary Tumor Markers for Bladder Cancer AHS – G2125

- Darwiche, F., Parekh, D. J., & Gonzalgo, M. L. (2015). Biomarkers for non-muscle invasive bladder cancer: Current tests and future promise. *Indian J Urol*, 31(4), 273-282. <https://doi.org/10.4103/0970-1591.166448>
- Davis, R., Jones, J. S., Barocas, D. A., Castle, E. P., Lang, E. K., Leveillee, R. J., Messing, E. M., Miller, S. D., Peterson, A. C., Turk, T. M., & Weitzel, W. (2012). Diagnosis, evaluation and follow-up of asymptomatic microhematuria (AMH) in adults: AUA guideline. *J Urol*, 188(6 Suppl), 2473-2481. <https://doi.org/10.1016/j.juro.2012.09.078>
- DeGeorge, K. C., Holt, H. R., & Hodges, S. C. (2017). Bladder Cancer: Diagnosis and Treatment. (1532-0650 (Electronic)). <https://www.aafp.org/pubs/afp/issues/2017/1015/p507.html>
- Dudley, J. C., Schroers-Martin, J., Lazzareschi, D. V., Shi, W. Y., Chen, S. B., Esfahani, M. S., Trivedi, D., Chabon, J. J., Chaudhuri, A. A., Stehr, H., Liu, C. L., Lim, H., Costa, H. A., Nabet, B. Y., Sin, M. L. Y., Liao, J. C., Alizadeh, A. A., & Diehn, M. (2019). Detection and Surveillance of Bladder Cancer Using Urine Tumor DNA. *Cancer Discov*, 9(4), 500-509. <https://doi.org/10.1158/2159-8290.Cd-18-0825>
- EAU. (2023). EAU Guidelines on Non-muscle Invasive Bladder Cancer. https://d56bochluxqnz.cloudfront.net/documents/full-guideline/EAU-Guidelines-on-Non-muscle-Invasive-Bladder-Cancer-2023_2023-03-10-101110_jued.pdf
- Ecke, T. H., Weiß, S., Stephan, C., Hallmann, S., Arndt, C., Barski, D., Otto, T., & Gerullis, H. (2018). UBC(®) Rapid Test-A Urinary Point-of-Care (POC) Assay for Diagnosis of Bladder Cancer with a focus on Non-Muscle Invasive High-Grade Tumors: Results of a Multicenter-Study. *Int J Mol Sci*, 19(12). <https://doi.org/10.3390/ijms19123841>
- Fantony, J. J., Longo, T. A., Gopalakrishna, A., Owusu, R., Lance, R. S., Foo, W. C., Inman, B. A., & Abern, M. R. (2017). Urinary NID2 and TWIST1 methylation to augment conventional urine cytology for the detection of bladder cancer. *Cancer Biomark*, 18(4), 381-387. <https://doi.org/10.3233/cbm-160261>
- FDA. (2018). Devices@FDA. Retrieved 11/12 from
- Grossman, H., Messing, E., Soloway, M., & et al. (2005). Detection of bladder cancer using a point-of-care proteomic assay. *JAMA*, 293(7), 810-816. <https://doi.org/10.1001/jama.293.7.810>
- Halpern, J. A., Chughtai, B., & Ghomrawi, H. (2017). Cost-effectiveness of Common Diagnostic Approaches for Evaluation of Asymptomatic Microscopic Hematuria. *JAMA Intern Med*, 177(6), 800-807. <https://doi.org/10.1001/jamainternmed.2017.0739>
- Hirasawa, Y., Pagano, I., Chen, R., Sun, Y., Dai, Y., Gupta, A., Tikhonenkov, S., Goodison, S., Rosser, C. J., & Furuya, H. (2021). Diagnostic performance of Oncuria™, a urinalysis test for bladder cancer. *Journal of Translational Medicine*, 19(1), 141. <https://doi.org/10.1186/s12967-021-02796-4>
- Hottinger, A. F., & Hormigo, A. (2011). Serum Biomarkers. In M. Schwab (Ed.), *Encyclopedia of Cancer* (pp. 3390-3394). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-16483-5_5269
- Kaufman, D. S., Shipley, W. U., & Feldman, A. S. (2009). Bladder cancer. *Lancet*, 374(9685), 239-249. [https://doi.org/10.1016/s0140-6736\(09\)60491-8](https://doi.org/10.1016/s0140-6736(09)60491-8)
- Landaas, E. J., Eckel, A. M., Wright, J. L., Baird, G. S., Hansen, R. N., & Sullivan, S. D. (2020). Application of Health Technology Assessment (HTA) to Evaluate New Laboratory Tests in a Health System: A Case Study of Bladder Cancer Testing. *Acad Pathol*, 7, 2374289520968225. <https://doi.org/10.1177/2374289520968225>

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Lerner, S. P., Raghavan, Derek. (2023, 12/18/2023). Overview of the initial approach and management of urothelial bladder cancer. <https://www.uptodate.com/contents/overview-of-the-initial-approach-and-management-of-urothelial-bladder-cancer>

Li, H. T., Duymich, C. E., Weisenberger, D. J., & Liang, G. (2016). Genetic and Epigenetic Alterations in Bladder Cancer. *Int Neurourol J*, 20(Suppl 2), S84-94. <https://doi.org/10.5213/inj.1632752.376>

Lopez-Beltran, A., Cheng, L., Gevaert, T., Blanca, A., Cimadamore, A., Santoni, M., Massari, F., Scarpelli, M., Raspollini, M. R., & Montironi, R. (2019). Current and emerging bladder cancer biomarkers with an emphasis on urine biomarkers. *Expert Rev Mol Diagn*, 1-13. <https://doi.org/10.1080/14737159.2020.1699791>

Lotan, Y., & Choueiri, T. (2022, 03/21/2022). Clinical presentation, diagnosis, and staging of bladder cancer. <https://www.uptodate.com/contents/clinical-presentation-diagnosis-and-staging-of-bladder-cancer>

Lotan, Y., Elias, K., Svatek, R. S., Bagrodia, A., Nuss, G., Moran, B., & Sagalowsky, A. I. (2009). Bladder cancer screening in a high risk asymptomatic population using a point of care urine based protein tumor marker. *J Urol*, 182(1), 52-57; discussion 58. <https://doi.org/10.1016/j.juro.2009.02.142>

Lotan, Y., & Roehrborn, C. G. (2003). Sensitivity and specificity of commonly available bladder tumor markers versus cytology: results of a comprehensive literature review and meta-analyses. *Urology*, 61(1), 109-118. [https://doi.org/10.1016/S0090-4295\(02\)02136-2](https://doi.org/10.1016/S0090-4295(02)02136-2)

Mahnert, B., Tauber, S., Kriegmair, M., Schmitt, U. M., Hasholzner, U., Reiter, W., Hofmann, K., Schmeller, N., & Stieber, P. (1999). BTA-TRAK--a useful diagnostic tool in urinary bladder cancer? *Anticancer Res*, 19(4a), 2615-2619. <https://pubmed.ncbi.nlm.nih.gov/10470204/>

Meleth, S., Reeder-Hayes, K., Ashok, M., Clark, R., Funkhouser, W., Wines, R., Hill, C., Shanahan, E., McClure, E., Burson, K., Coker-Schwimmer, M., Garge, N., & Jonas, D. E. (2014). AHRQ Technology Assessments. In *Technology Assessment of Molecular Pathology Testing for the Estimation of Prognosis for Common Cancers*. Agency for Healthcare Research and Quality (US). https://www.ncbi.nlm.nih.gov/books/NBK285410/pdf/Bookshelf_NBK285410.pdf

Mitra, A., Birkman, M., Penson, D., & Cote, R. (2023, 12/18/2023). Urine biomarkers for the detection of urothelial (transitional cell) carcinoma of the bladder <https://www.uptodate.com/contents/urine-biomarkers-for-the-detection-of-urothelial-transitional-cell-carcinoma-of-the-bladder>

Monteiro, L. L., Witjes, J. A., Agarwal, P. K., Anderson, C. B., Bivalacqua, T. J., Bochner, B. H., Boormans, J. L., Chang, S. S., Domínguez-Escrig, J. L., & McKiernan, J. M. J. W. j. o. u. (2018). ICUD-SIU International Consultation on Bladder Cancer 2017: management of non-muscle invasive bladder cancer. 1-10. <http://urology.stanford.edu/content/dam/sm/urology/JJimages/publications/ICUD-SIU-International-Consultation-on-Bladder-Cancer-2017-management-of-non-muscle-invasive-bladder-cancer.pdf>

Mossanen, M., Wang, Y., Szymaniak, J., Tan, W. S., Huynh, M. J., Preston, M. A., Trinh, Q. D., Sonpavde, G., Kibel, A. S., & Chang, S. L. (2019). Evaluating the cost of surveillance for non-muscle-invasive bladder cancer: an analysis based on risk categories. *World J Urol*, 37(10), 2059-2065. <https://doi.org/10.1007/s00345-018-2550-x>

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Moyer, V. A. (2011). Screening for bladder cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*, 155(4), 246-251. <https://doi.org/10.7326/0003-4819-155-4-201108160-00008>

NCCN. (2023a). Bladder Cancer - Version 3.2023
https://www.nccn.org/professionals/physician_gls/pdf/bladder.pdf

NCCN. (2023b). NCCN Practice Guidelines: Bladder Cancer Version 3.2023.
https://www.nccn.org/professionals/physician_gls/pdf/bladder.pdf
NCI. (2022). Bladder and Other Urothelial Cancers Screening (PDQ®)—Health Professional Version. U.S. Department of Health and Human Services. Retrieved 01/29/2023 from
<https://www.cancer.gov/types/bladder/hp/bladder-screening-pdq>

NICE. (2023). URO17 for detecting bladder cancer.
<https://www.nice.org.uk/advice/mib250/chapter/The-technology>

Nonagen Bioscience. (2022). Bladder Cancer. <https://www.nonagen.com/products>

Nucleix. (2015). BLADDER EPICHECK <https://www.nucleix.com/bladder-epicheck/>

Pangea. (2019). Pangea® Laboratory to License Bladder CARE™ Technology from Zymo Research. <https://www.prnewswire.com/news-releases/pangea-laboratory-to-license-bladder-care-technology-from-zymo-research-300800622.html>

Pangea. (2020). Features of Bladder CARE™. <https://www.pangealab.com/bladdercare/>

Perazalla, M. (2021, 07/09/2021). Etiology and evaluation of hematuria in adults.
<https://www.uptodate.com/contents/etiology-and-evaluation-of-hematuria-in-adults>

Piao, X. M., Jeong, P., Kim, Y. H., Byun, Y. J., Xu, Y., Kang, H. W., Ha, Y. S., Kim, W. T., Lee, J. Y., Woo, S. H., Kwon, T. G., Kim, I. Y., Moon, S. K., Choi, Y. H., Cha, E. J., Yun, S. J., & Kim, W. J. (2019). Urinary cell-free microRNA biomarker could discriminate bladder cancer from benign hematuria. *Int J Cancer*, 144(2), 380-388. <https://doi.org/10.1002/ijc.31849>

Pichler, R., Fritz, J., Tulchiner, G., Klinglmair, G., Soleiman, A., Horninger, W., Klocker, H., & Heidegger, I. (2018). Increased accuracy of a novel mRNA-based urine test for bladder cancer surveillance. *BJU Int*, 121(1), 29-37. <https://doi.org/10.1111/bju.14019>

Protean Biodiagnostics. (2021). URO17 is the Most Sensitive and Specific Urine Test for Bladder Cancer. <https://www.proteanbiodx.com/uro17>

Quest. (2020). Bladder Tumor Antigen DetectR™.
<https://testdirectory.questdiagnostics.com/test/test-detail/34055/bladder-tumor-antigen-detectr?cc=MASTER>

Sathianathan, N. J., Butaney, M., Weight, C. J., Kumar, R., & Konety, B. R. (2018). Urinary Biomarkers in the Evaluation of Primary Hematuria: A Systematic Review and Meta-Analysis. *Bladder Cancer*, 4(4), 353-363. <https://doi.org/10.3233/blc-180179>

Schmitz-Dräger, B. J., Droller, M., Lokeshwar, V. B., Lotan, Y., Hudson, M. A., van Rhijn, B. W., Marberger, M. J., Fradet, Y., Hemstreet, G. P., Malmstrom, P. U., Ogawa, O., Karakiewicz, P. I., & Shariat, S. F. (2015). Molecular Markers for Bladder Cancer Screening, Early Diagnosis, and Surveillance: The WHO/ICUD Consensus. *Urologia Internationalis*, 94(1), 1-24.
<https://doi.org/10.1159/000369357>

Urinary Tumor Markers for Bladder Cancer AHS – G2125

Soubra, A., & Risk, M. C. (2015). Diagnostics techniques in nonmuscle invasive bladder cancer. *Indian J Urol*, 31(4), 283-288. <https://doi.org/10.4103/0970-1591.166449>

Sutton, A. J., Lamont, J. V., Evans, R. M., Williamson, K., O'Rourke, D., Duggan, B., Sagoo, G. S., Reid, C. N., & Ruddock, M. W. (2018). An early analysis of the cost-effectiveness of a diagnostic classifier for risk stratification of haematuria patients (DCRSHP) compared to flexible cystoscopy in the diagnosis of bladder cancer. *PLoS One*, 13(8), e0202796. <https://doi.org/10.1371/journal.pone.0202796>

Tan, W. S., Tan, W. P., Tan, M. Y., Khetrupal, P., Dong, L., deWinter, P., Feber, A., & Kelly, J. D. (2018). Novel urinary biomarkers for the detection of bladder cancer: A systematic review. *Cancer Treat Rev*, 69, 39-52. <https://doi.org/10.1016/j.ctrv.2018.05.012>

USPSTF. (2021). Bladder Cancer in Adults: Screening. <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/bladder-cancer-in-adults-screening>

Vasdev, N., Hampson, A., Agarwal, S., Swamy, R., Chilvers, M., Hampson, A., Jahanfard, S., & Kim, N. (2021). The role of URO17™ biomarker to enhance diagnosis of urothelial cancer in new hematuria patients—First European Data. *BJUI Compass*, 2(1), 46-52. <https://doi.org/10.1002/bco2.50>

Witjes, J. A., Bruins, H. M., Cathomas, R., Compérat, E. M., Cowan, N. C., Gakis, G., Hernández, V., Linares Espinós, E., Lorch, A., Neuzillet, Y., Rouanne, M., Thalmann, G. N., Veskimäe, E., Ribal, M. J., & van der Heijden, A. G. (2022). Muscle-invasive and Metastatic Bladder Cancer. <https://uroweb.org/guideline/bladder-cancer-muscle-invasive-and-metastatic/>

Zuiverloon, T. C. M., de Jong, F. C., & Theodorescu, D. (2017). Clinical Decision Making in Surveillance of Non-Muscle-Invasive Bladder Cancer: The Evolving Roles of Urinary Cytology and Molecular Markers. *Oncology (Williston Park)*, 31(12), 855-862. <https://www.cancernetwork.com/view/clinical-decision-making-surveillance-nonmuscle-invasive-bladder-cancer-evolving-roles-urinary>

Policy Implementation/Update Information

- 1/1/19 New policy developed. BCBSNC will provide coverage for urinary tumor markers for bladder cancer when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (sk)
- 10/1/19 Policy statement revised to read: BCBSNC will provide coverage for Urinary Tumor Markers for Bladder Cancer when it is determined the medical criteria or reimbursement guidelines below are met. Wording revised in the When Covered section. "Medically necessary" changed to read "Reimbursement is allowed..." Wording revised in the Not Covered section. "Investigational" changed to read "Reimbursement is not allowed..." Deleted coding grid. Notification given 10/1/2019 for effective date 12/2/2019.
- 10/15/19 Medical Director review. Statement added that reimbursement for the use of all other urinary biomarkers, including but not limited to quantitative mRNA (Cxbladder), is not allowed. Notification given 10/15/2019 for effective date 12/31/2019. (sk)
- 12/31/19 Specialty Matched Consultant Advisory Panel review 11/20/2019. (sk)

Urinary Tumor Markers for Bladder Cancer AHS – G2125

- 5/12/20 Reviewed by Avalon 1st Quarter 2020 CAB. Medical Director review 4/2020. References updated and added. Description section updated. Related Policies added. When Covered section updated. When Not Covered section updated. State and Federal Regulations section updated. Policy Guidelines updated. Codes 88271, 88299, and 88365 deleted from Billing/Coding section. Codes 0012M and 0013M added to Billing/Coding section. (sk)
- 5/4/21 Specialty Matched Consultant Advisory Panel review 11/18/2020. Reviewed by Avalon 1st Quarter 2021 CAB. Medical Director review 4/2021. References updated and added. Description section updated. Related Policies updated. State and Federal Regulations section updated. Policy Guidelines updated. (sk)
- 12/14/21 Specialty Matched Consultant Advisory Panel review 11/17/2021. (sk).
- 5/17/22 Reviewed by Avalon 1st Quarter 2022 CAB. Medical Director review 4/2022. References updated and added. Description section updated. Removed the following statement from the When Not Covered section; “The use of all other urinary biomarkers, including but not limited to quantitative mRNA (Cxbladder) is investigational.” Policy Guidelines updated. (sk)
- 9/13/22 Reviewed by Avalon 2nd Quarter 2022 CAB. Medical Director review 8/2022. Related policy “Molecular Panel Testing of Cancers for Diagnosis, Prognosis, and Identification of Targeted Therapy” removed. Table of Terminology added. References updated and added. Description section updated. Added codes 88346 and 88350 to Billing/Coding section. Policy Guidelines updated. (sk)
- 12/30/22 Added code 0363U to Billing/Coding section. (sk)
- 4/18/23 Added new codes 0365U, 0366U, 0367U to Billing/Coding section for effective date 4/1/2023. (sk)
- 5/16/23 Reviewed by Avalon 1st Quarter 2023 CAB. Medical Director review 4/2023. Related References updated and added. Description section updated. Policy Guidelines updated. (sk)
- 5/15/24 Reviewed by Avalon 1st Quarter 2024 CAB. Updated the background, guidelines and recommendations. Updated table of terminology and references. Medical Director review. No changes in coverage criteria. Added CPT code 0420U to the billing section. (rp)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.